

**Table 3. PPOR Model - Comparison of Excess Fetal Infant Mortality Rates (FIMR)<sup>a</sup> in Kansas, 2005-2009**

Group	Maternal Health/ Prematurity		Maternal Care		Newborn Care		Infant Health		Total	
	Groups Specific Rate	Excess Rate <sup>c</sup>	Groups Specific Rate	Excess Rate <sup>c</sup>	Groups Specific Rate	Excess Rate <sup>c</sup>	Groups Specific Rate	Excess Rate <sup>c</sup>	Groups Specific Rate	Excess Rate <sup>c</sup>
U.S. Reference Group <sup>b</sup>	2.2		1.5		1.1		0.9		5.7	
Kansas	3.1	0.9	2.1	0.6	1.5	0.4	2.3	1.4	9.0	3.3
		186 deaths		124 deaths		83 deaths		289 deaths		681 deaths
White non- Hispanic	2.7	0.5	1.9	0.4	1.5	0.4	2.0	1.1	8.1	2.4
		74 deaths		59 deaths		59 deaths		163 deaths		356 deaths
Black non- Hispanic	7.2	5.0	3.8	2.3	1.7	0.6	3.9	3.0	16.6	10.9
		71 deaths		33 deaths		9 deaths		43 deaths		155 deaths
Hispanic	3.2	1.0	2.5	1.0	1.3	0.2	2.6	1.7	9.6	3.9
		33 deaths		33 deaths		7 deaths		56 deaths		129 deaths

Source: Bureau of Epidemiology and Public Health Informatics, KDHE

<sup>a</sup> Fetal and infant mortality per 1,000 live births (plus fetal deaths)

<sup>b</sup> The USA reference group concerns fetal infant deaths to white non-Hispanic mothers greater than 19 years of age, with at least 13 years of education, 2000-2002 combined.

<sup>c</sup> Excess rate is calculated by the group specific rate minus the USA reference rate.